

Colonoscopy – Patient Instructions

For appointments at or after 11 a.m.

This sheet is double sided. Please read and follow all instructions carefully to ensure the highest level of safety and effectiveness. You must have someone to accompany you home; you cannot drive.

Medications:

Please make sure all of your medications are cleared by a doctor.

Stop 7 days prior	Aspirin(ASA 81 mg, Entrophen) Iron Tablets Xarelto	
Stop 5 days prior	Coumadin Pradaxa	
Stop 3 days prior	Ticlopidine (Ticlid) Clopidogrel (Plavix) Aggrenox	
Special instructions for diabetics	Day before	Day of
Hypoglycemic medications (tablets)	Take ½ usual dose	None (Until after procedure with food)
Insulin	Take ½ usual dose	None (Until after procedure with food)

Continue taking **blood pressure medication** as normal even on the day of your procedure, with a small sip of water. Do not drink any more than a small sip. It is important that your blood pressure remains under control.

Colonoscopy Preparation Checklist

Confirm your medications: Speak with your doctor about any medication that you may need to stop taking before your test.

Follow the preparation: Please carefully read all of the instructions provided on both sides of this sheet, simply complete the steps and follow the dietary restrictions before your test.

Watch the foods you eat: Follow the examples provided about what you can and cannot eat before your test.

Dress comfortably: Upon entering the procedure room you will be provided with a gown to change into for the test, we recommend wearing something easy to change in and out of.

Do not consume any cannabis related products for 48 hours prior to your procedure.

Confirm your ride home: Because of the sedatives used during colonoscopy, you will need someone to drive you home or accompany you in a taxi. The accompanying adult **MUST** sign you out from the recovery area. You may not leave alone. If you do not bring someone your procedure will be cancelled.

7 Days Prior to Colonoscopy

Purchase one PegLyte kit and Dulcolax 3 x 5mg tablets

- *Please note that substituting PegLyte with a generic alternative may compromise the effectiveness of your colon preparation.*
-

5 Days Prior to Colonoscopy

Stop eating foods that contain **nuts/seeds** (they can damage the scopes).

1 Day Prior to Colonoscopy

No foods allowed. Drink clear liquids only (at least one glass every hour) up until 3 hours prior to your colonoscopy. If you are also having a gastroscopy, drink clear liquids only up until 3 hours prior to the procedure.

Examples of clear liquids you may consume: Water, apple juice, white grape juice, white cranberry juice, Gatorade®, Powerade®, Gastrolyte®, soda water, jello, and clear soup(broth only). **NO RED/PURPLE DRINKS OR ALCOHOL.** Coffee or tea you may add sugar or sweetener. **NO CREAM/MILK.** Remember: If you can see through it, you can drink it.

Make sure you have access to a washroom; the laxative will induce bowel movements.

Morning – Mix Peglyte powder into 4 litres of cold water (fill to the marked line on the bottle) and mix rapidly until fully dissolved. Refrigerate until ready to drink in the evening.

At **3:00 p.m.** take 3 dulcolax tablets. This medication only softens the stool.

At **7:00 p.m.** Drink 2 litres (half) of the PegLyte preparation. You must consume one cup (250ml) every 15 minutes over the following 2 hours. Refrigerate remainder. Continue to drink clear fluids.

Day of Colonoscopy

No foods allowed until 1 hour after your procedure.

5 hours before appointment Drink 2 litres (half) of the PegLyte preparation. You must consume one cup (250ml) every 15 minutes over the following 2 hours.

You may only drink water on the day of your procedure until 3 hours before your procedure time.

STOP ALL LIQUIDS 3 HOURS BEFORE YOUR TEST. You must have a completely empty stomach prior to your test as even small amounts of food/liquid may result in delay or cancellation and increase the risk of aspiration pneumonia that could result in death.

Bring your health card. Ensure you have someone to drive you home or accompany you in a taxi.

For more information about the procedure including educational literature and videos, please access our website www.gtaendoscopy.com.

Please note that there is a **cancellation fee of \$100** if you cancel less than 3 business days before your procedure without a medical note.

Location – We are located on the southeast corner of Markham Road and Milner Avenue.

GTA Endoscopy Services: 305 Milner Avenue, Suite 920. Scarborough, ON.

Phone: For any questions or concerns we may be reached at (416) 321-3883.